

List of studies

Coy JF: EDIM-TKTL1/Apo10 Blood Test: An Innate Immune System Based Liquid Biopsy for the Early Detection, Characterization and Targeted Treatment of Cancer. *Int J Mol Sci.* 2017 Apr 20;18(4). pii: E878. doi: 10.3390/ijms18040878. Zyagnum Frankfurt, Germany.

Xu. W.; Lai. R.K.; Lin. S.H.; Tse, A.P.; Chiu. D.K.; Koh. H.Y.; Law. C.T.; Wong. C.M.; Cai. Z.; Wong. C.C.: et al. Transketolase counteracts oxidative stress to drive cancer development. *Proc Natl Acad Sci U S A.* 2016 Feb 9;113(6):E725-34. The University of Hong Kong. Hong Kong. China;

Li, B.; Iglesias-Pedraz, J.M.; Chen, L.Y.; Yin, F.; Cadenas, E.; Reddy, S.; Comai, L. Downregulation of the Werner syndrome protein induces a metabolic shift that compromises redox homeostasis and limits proliferation of cancer cells. *Aging Cell* 2014, Apr. 13, 367-378. University of Southern California, Los Angeles. USA / University of Southern California, Los Angeles, USA

Jayachandran, A.; Lo. P.H.; Chueh. A.C.; Rrithviraj, P.; Molania. R.; Davalos-Salas, M.; Anaka. M.; Walkiewicz, M.; Cebon, J.; Behren, A. Transketolase-like 1 ectopic expression is associated with DNA hypomethylation and induces the Warburg effect in melanoma cells. *BMC Cancer.* 2016 Feb 22;16:134. Ludwig Institute for Cancer Research, Heidelberg. Australia.

Li, J.; Zhu, S.C.; Li. S.G.; Zhao, Y.; Xu, J.R.; Song, C.Y. TKTL1 promotes cell proliferation and metastasis in esophageal squamous cell carcinoma. *Blamed Pharmacother.* 2015 Aug;74:71-6. The Fourth Hospital of Hebei Medical University, Shijiazhuang, China.

Zerilli Met al: Increased expression of transketolase-like-1 in papillary thyroid carcinomas smaller than 1.5 cm in diameter is associated with lymph-node metastases. *Cancer.* 2008 Sep 1;113(5):936-44. University of Palermo, Italy.

Ahopelto, K.; Bockelman, C.; Hagstrom, J.; Koskensalo, S.; Haglund. C. Transketolase-like protein 1 expression predicts poor prognosis in colorectal cancer. *Cancer Biol. Ther.* 2016, 17, 163-168. University of Helsinki and Helsinki University Hospital, Helsinki, Finland.

Bentz Set al: Hypoxia induces the expression of transketolase-like 1 in human colorectal cancer. *Digestion.* 2013;88(3):182-92. University Hospital Zurich, Switzerland.

Jansen N and Coy JF: Diagnostic use of EDIM-blood test for early detection of colon cancer metastasis. *Future Oncol.* 2013;9(4):605-9.

Schwaab Jet al: Expression of Transketolase like gene 1 (TKTL1) predicts disease-free survival in patients with locally advanced rectal cancer receiving neoadjuvant chemoradiotherapy. *BMC Cancer* 2011;11:363. University Medical Centre Mannheim, Germany.

Diaz-Morali Set al: Transketolase-like 1 expression is modulated during colorectal cancer progression and metastasis formation. *PLoS One* 2011;6(9):e25323. Epub 2011 Sep 27. University of Barcelona. Spain.

Xu X et al. Transketolase-like protein 1 (TKL1) is required for rapid cell growth and full viability of human tumor cells, *Int J Cancer*, 2009 Mar 15;124(6):1330-7. DKFZ German Cancer Research Centre, Heidelberg, Germany.

Langbein S et al: Expression of transketolase TKTL1 predicts colon and urothelial cancer patient survival: Warburg effect reinterpreted. *Br J Cancer*. 2006 Feb 27;94(4):578-85. University Hospital Mannheim, Germany.

Dong, Y.; Wang, M. Knockdown of TKTL1 additively complements cisplatin-induced cytotoxicity in nasopharyngeal carcinoma cells by regulating the levels of NADPH and ribose-5-phosphate. *Biomed. Pharmacother*. 2017, 85, 672— 678. Central Hospital Affiliated to Zhengzhou University, Zhengzhou, China.

Song, Y.; Liu, D.; He, G. TITLfi and p63 are biomarkers for the poor prognosis of gastric cancer patients. *Cancer Biomark*. 2015, 5, 591—597. Central South University, Changsha, China.

Yuan W et al: Silencing of TKTL1 by siRNA inhibits proliferation of human gastric cancer cells in vitro and in vivo. *Cancer Biol Ther*. 2010 May;9(9):710-6. Epub 2010 May. Central South University, Changsha, Hunan, China.

Staiger W et al: Expression of the mutated transketolase TKTL1, a molecular marker in gastric cancer. *Oncol Rep*. 2006 Oct;16(4):657-61. Universitätsmedizin Mannheim. University Hospital Mannheim, Germany. Rotmann A et al: A new diagnostic approach for the early detection and monitoring of breast cancer patients. Poster Glasgow 9th European Breast Cancer Conference (EBCC-9), 19 to 21 March 2014 Rotmann A et al: A new diagnostic test for monitoring of breast cancer patients. Poster, Deutscher Krebskongress 2014

Grimm M et al: A biomarker based detection and characterization of carcinomas exploiting two fundamental biophysical mechanisms in mammalian cells. *BMC Cancer* 2013, 13:569. Tuebingen University Hospital, DKFZ German Cancer Research Center, Heidelberg, Germany.

Rotmann A: The use of new diagnostic tests for the monitoring of new and existing therapies for breast cancer patients. *J. Clin. Oncol.*, ASCO Meeting Abstracts. 2013; 31:e22006

Feyen O et al: EDIM-TKTL1 blood test: a noninvasive method to detect upregulated glucose metabolism in patients with malignancies. *Future Oncol*. 2012;8(10):1349-59. Central Clinic Bad Berka, Germany. Feyen O et al: EDIM blood test: a non-invasive method to detect patients with malignancies. *Cell Symposia* July 6-8, 2012: Poster

Rotmann A et al: Apo10 - a new biomarker for early detection of disorders of cell proliferation and solid tumours. *Int J Gynaecol Obstet* 119, Supplement 3 (2012), 5466. Abstract FIGO

Schmidt M et al: Glycolytic phenotype in breast cancer: activation of Akt, up-regulation of GLUT1, TKTL1 and downregulation of M2PK. *J Cancer Res Clin Oncol*. 2010 Feb;136(2):219-25. Epub 2009 Aug 5. University of Wuerzburg, Germany.

Rotmann A et al: TKTL1 - a new biomarker and its relevance in the daily gynaecological practice. Poster FIGO 2009

Foeldi M et al: Transketolase protein TKTL1 overexpression: A potential biomarker and therapeutic target in breast cancer. *Oncol Rep*. 2007 Apr;17(4):841-5. University Hospital Freiburg, Germany. Krockenberger M et al: Expression of transketolase-like 1 protein (TKTL1) in human endometrial cancer. *Anticancer Res*. 2010 May;30(5):1653-9. University of Wuerzburg, Germany.

Ricciardelli, C.; Lokman, N.A.; Cheruvu, S.; Tan, I.A.; Ween, M.P.; Pyragius, C.E.; Ruszkiewicz, A.; Hoffmann, P.; Oehler, M.K. Transketolase is upregulated in metastatic peritoneal implants and promotes ovarian cancer cell proliferation. *Clin. Exp. Clin Exp*

Metastasis. 2015 Jun;32(5):441-55. University of Adelaide, Adelaide, Australia

Schmidt M et al: Glucose metabolism and angiogenesis in granulosa ocell tumors of the ovary: activation of Akt, expression of M2PK, TKTL1 and VEGF. *Eur J Obstet Gynecol Reprod Biol.* 2008 Jul;139(1):72-8. Epub 2008 Apr 3. University of Wuerzburg, Germany.

Chen H et al: Overexpression of transketolase-like gene 1 is associated with cell proliferation in uterine cervix cancer. *J Exp Clin Cancer Res.* 2009 Mar 30;28:43.

Kohrenhagen N et al: Expression of transketolase-like 1 (TKTL1) and p-Akt correlates with the progression of cervical neoplasia. *J Obstet Gynaecol Res.* 2008 Jun;34(3):293-300. University of Wuerzburg, Wuerzburg, Germany. Wanka C et al: Tp53-induced glycolysis and apoptosis regulator (TIGAR) protects glioma cells from starvation-induced cell death by upregulating respiration and improving cellular redox homeostasis. *J Biol Chem.* 2012 Sep 28;287(40):33436-46. Epub 2012 Aug 10. Goethe University Frankfurt, Germany.

Wani K et al: A prognostic gene expression signature in infratentorial ependymoma. For the Collaborative Ependymoma Research Network. *Acta Neuropathol.* 2012 May;123(5):727-38. Epub 2012 Feb 10. The University of Texas, MD Anderson Cancer Center, Houston, Texas, USA.

Völker HU et al: Expression of transketolase-like 1 and activation of Akt in grade IV glioblastomas compared with grades II and III astrocytic gliomas. *Am J Clin Pathol.* 2008 Jul;130(1):50-7. University of Wuerzburg, Germany.

Lange CA et al: Enhanced TKTL1 Expression in Malignant Tumors of the Ocular Adnexa Predicts Clinical Outcome. *Ophthalmology.* 2012 Sep;119(9):1924-9. Epub 2012 Jun 1. University Hospital Freiburg, Germany.

Dong, Y.; Wang, M. Knockdown of TKTL1 additively complements cisplatin-induced cytotoxicity in nasopharyngeal carcinoma cells by regulating the levels of NADPH and ribose-5-phosphate. *Biomed. Pharmacother.* 2017, 85, 672— 678. Zhengzhou University, Zhengzhou, China.

Grimm, M.; Hoefert, S.; Krimmel, M.; Biegner, T.; Feyen, O.; Teriete, P.; Reinert, S. Monitoring carcinogenesis in a case of oral squamous cell carcinoma using a panel of new metabolic blood biomarkers as liquid biopsies. *Oral Maxillofac Surg.* 2016 Sep;20(3):295-302. Tuebingen University Hospital, Tuebingen, Germany

Diaz-Moralli, S.; Aguilar, E.; Marin, S.; Coy, J.F.; Dewerchin, M.; Antoniewicz, M.R.; Meca-Cortés, O.; Notebaert, L.; Ghesquiére, B.; Eelen, G.; et al. A key role for Transketolase-like 1 in tumor metabolic reprogramming. *Oncotarget.* 2016 Aug 9;7(32):51875-51897. Universitat de Barcelona, Barcelona, Spain

Grimm, M.; Feyen, O. ; Coy, J.F.; Hofmann, H.; Teriete, P.; Reinert, S. Analysis of circulating CD14+/CD16+ monocytederived macrophages (MDMs) in the peripheral blood of patients with oral squamous cell carcinoma. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2016 Mar;121(3):301-6. Tuebingen University Hospital, Tuebingen, Germany / Zyagnum AG Frankfurt, Germany

Grimm, M.; Kraut, W.; Hoefert, S.; Krimmel, M.; Biegner, T.; Teriete, P.; Cetindis, M.; Polligkeit, J.; Kluba, S.; Munz, A.; et al. Evaluation of a biomarker based blood test for monitoring surgical resection of oral squamous cell carcinomas. *Clin. Oral Investig.* 2016, 20, 329—338. University Hospital Tuebingen, Germany / Sanford-Burnham Medical Research InstituteLaJollaUSA

Grimm et al: Apoptosis resistance-related ABCBs and DNaseX (Apo10) expression in oral carcinogenesis. *Acta Odontol Scand*. 2014 Sep 19;1-7. (Epub ahead of print). Tuebingen University Hospital, Tuebingen, Germany.

Grimm et al.: Association of cancer metabolism-related proteins with oral carcinogenesis — indications for chemoprevention and metabolic sensitizing of oral squamous cell carcinoma?. *J Transl Med*. 2014 Jul 21;12:208. Tuebingen University Hospital, Tuebingen, Germany.

Grimm et al: GLUT-1+/TKTL1+ coexpression predicts poor outcome in oral squamous cell carcinoma. *Oral Surg Oral Med Oral Pathol Oral Radial*. 2014 Jun;117(6):743-53. Tuebingen University Hospital, Tuebingen,

Grimm M et al: A biomarker based detection and characterization of carcinomas exploiting two fundamental biophysical mechanisms in mammalian cells. *BMC Cancer*. 2013, 13:569. Tuebingen University Hospital, Tuebingen, DKFZ German Cancer Research Center, Heidelberg, Germany.

Grimm M et al: Transketolase-like protein 1 expression in recurrent oral squamous cell carcinoma after curative resection: a case report. *Oral Surg Oral Med Oral Pathol Oral Radial*. 2012 Jul 20. Tuebingen University Hospital, Tuebingen, Germany.

Hartmannsberger D et al: Transketolase-like protein 1 confers resistance to serum withdrawal in vitro. *Cancer Lett*. 2011 Jan 1;300(1):20-9. Epub 2010 Sep 29. Ludwig-Maximilians-University of Munich, Germany.

Sun W et al: TKTL1 is activated by promoter hypomethylation and contributes to head and neck squamous cell carcinoma carcinogenesis through increased aerobic glycolysis and HIF1alpha stabilization. *Clin Cancer Res*. 2010 Feb 1;16(3):857-66. Epub 2010 Jan 26

Smith IM et al: Coordinated activation of candidate proto-oncogenes and cancer testes antigens via promoter demethylation in head and neck cancer and lung cancer. *PLoS One*. 2009;4(3):e4961. Epub 2009 Mar 23

Völker HU et al: Overexpression of transketolase TKTL1 is associated with shorter survival in laryngeal squamous cell carcinomas. *Eur Arch Otorhinolaryngol*. 2007 Dec;264(12):1431-6. Epub 2007 Jul 18. University of Wuerzburg, Germany.

Feyen O et al: EDIM-TKTL1 blood test: a noninvasive method to detect upregulated glucose metabolism in patients with malignancies. *Future Oncol*. 2012;8(10):1349-59. Central Clinic Bad Berka, Germany. Feyen O et al: EDIM blood test: a non-invasive method to detect patients with malignancies. *Cell Symposia* July 6-8, 2012 Poster

Kayser G et al: Poor outcome in primary non-small cell lung cancers is predicted by transketolase TKTL1 expression. *Pathology*. 2011 Dec;43(7):719-24. University Hospital Freiburg, Germany.

Kayser G et al: Lactate-dehydrogenase 5 is overexpressed in non-small cell lung cancer and correlates with the expression of the transketolase-like protein 1. *Diagn Pathol*. 2010 Apr 12;5:22. University Hospital Freiburg, Germany.

Prasad V et al: Transketolase-like 1 (TKTL1), a new intracellular marker for staging of non-small cell lung cancer (NSCLC): Correlation with metabolic tumor size on F-18 FDG PET/CT. *Poster SNM 2010*. Central Clinic Bad Berka, Germany

Smith IM et al: Coordinated activation of candidate proto-oncogenes and cancer testes

antigens via promoter demethylation in head and neck cancer and lung cancer. PLoS One. 2009;4(3):e4961. Epub 2009 Mar 23

Grimm M et al: A biomarker based detection and characterization of carcinomas exploiting two fundamental biophysical mechanisms in mammalian cells. BMC Cancer.2013, 13:569. Tuebingen University Hospital, DKFZ German Cancer Research Center, Heidelberg, Germany.

Arnoldt J: Therapy monitoring and early detection of metastasis using tumor protein detection in macrophages. Anticancer Research ISSN 0250-7005: 2011 May:2012

Langbein S et al: Expression of transketolase TKTL1 predicts colon and urothelial cancer patient survival: Warburg effect reinterpreted. Br J Cancer. 2006 Feb 27;94(4):578-85. University Hospital Mannheim, Germany.

Wu HT et al: Anaplastic neuroblastomas express transketolase-like enzyme 1. J Clin Pathol. 2009 May;62(5):460-3.Epub 2009 Jan 12

Langbein S et al: Metastasis is promoted by a bioenergetic switch: new targets for progressive renal cell cancer. Int J Cancer. 2008 Jun 1;122(11):2422-8. University of Amsterdam, The Netherlands.

2018-05 / pictures: Shutterstock: angellodeco | Alex Traksel | Have a nice day Photo